For loss of use insurance, the following X-ray studies are required for assessment:

Horseshoes must be removed for X-ray examination!

All X-ray studies must be clearly marked with the name of the horse, date, side markings and be in DICOM format.

<u>Toe - forelimb (central view radial leg):</u>

- lateromedial (LM)
- dorsoproximal palmarodistal oblique (Oxspring)
- palmaroproximal palmarodistal oblique (Morgan, Skyline)

Fetlock – front/hind limbs:

- lateromedial (LM)
- dorsopalmar (DP)
- dorsomedial palmaro-/plantarolateral oblique (DMPL)
- dorsolateral palmaro-/plantaromedial oblique (DLPM)

Hock:

- dorsoplantar (DP) incl. prox. Metatarsus
- lateromedial (LM)
- dorsomedial plantarolateral oblique (DMPL)
- dorsolateral plantaromedial oblique (DLPM)

Stifle joint:

- caudocranial (CdCr) or caudolateral craniomedial oblique (CdL-CrM)
- lateromedial (LM)

Ultrasound:

Transversal and horizontal views; one image per zone, suspensory ligament branches individually

Metacarpus/-tarsus bds./ Flexor tendon apparatus

• superficial, deep flexor tendons and suspensory ligament branches

Furthermore, images / views of regions that have been noticed by the clinical and radiological examination with deviating findings.